



UNIVERSIDADE FEDERAL DO CEARÁ  
CENTRO DE CIÊNCIAS AGRÁRIAS  
DEPARTAMENTO DE ZOOTECNIA  
PROGRAMA DE PÓS-GRADUAÇÃO EM ZOOTECNIA  
FORRAGICULTURA – AFP740

LITERATURA SUGERIDA PARA A PARTE I (Prof. Magno J. D. Cândido)

TERMINOLOGIA

- FORAGE AND GRAZING TERMINOLOGY COMMITTEE. Terminology for grazing lands and grazing animals. **J. Prod. Agric.**, v.5, n.1, p.191-201, 1992.
- HODGSON, J. Nomenclature and definitions in grazing studies. **Grass Forage Sci.**, v.34, n.1, p.11-18, 1979.
- THOMAS, H. Terminology and definitions in studies of grassland plants. **Grass Forage Sci.**, v.36, n.2, p.97-105, 1981.

ALONGAMENTO FOLIAR

- GASTAL, F. et al. A model of leaf extension rate of Tall Fescue in response to nitrogen and temperature. **Ann. Bot.**, v.70, n.5, p.437-442, 1992.
- KRIEDEMANN. Stomatal and photosynthetic limitations to leaf growth. **Aust. J. Plant Physiol.**, v.13, p.15-31, 1986.
- PEACOCK, J.M. Temperature and leaf growth in *Lolium perenne*. I. The thermal microclimate: its measurement and relation to crop growth. **J. Appl. Ecol.**, v.12, n.1, p.99-113, 1975.
- PEACOCK, J.M. Temperature and leaf growth in *Lolium perenne*. II. The site of temperature perception. **J. Appl. Ecol.**, v.12, n.1, p.115-123, 1975.
- RYLE, G.J.A. A comparison of leaf and tiller growth in seven perennial grasses as influenced by nitrogen and temperature. **J. Br. Grassl. Soc.**, v.19, n.3, p.281-290, 1964.
- SILSBURY, J.H. Leaf growth in pasture grasses. **Trop. Grassl.**, v.4, n.1, p.17-36, 1970.
- SILSBURY, J.H. The effects of temperature and light energy on dry weight and leaf area changes in seedling plants of *Lolium perenne* L. **Austr. J. Agric. Res.**, v.22, n.2, p.177-187, 1971.
- SKINNER, R.H. e NELSON, C.J.. Epidermal cell division and the coordination of leaf and tiller development. **Ann. Bot.**, v.74, n.1, p.9-15, 1994.
- SKINNER e NELSON. Elongation of the grass leaf and its relationship to the phyllochron. **Crop Sci.**, v.35, n.1, p.4-10, 1995.

APARECIMENTO FOLIAR

- ANSLOW, R.C. The rate of appearance of leaves on tillers of the gramineae. **Herbage Abst.**, v.36, n.3, p.149-155, 1966.
- HIRATA, M. Effects of nitrogen fertiliser rate and cutting height on leaf appearance and extension in bahia grass (*Paspalum notatum*) swards. **Trop. Grassl.**, v.34, p.7-13, 2000.
- SKINNER, R.H., NELSON, C.J. Role of leaf appearance rate and the coleoptile tiller in regulating tiller production. **Crop Sci.**, v.34, n.1, p.71-75, 1994.
- Van ESBROECK, G.A., HUSSEY, M.A., SANDERSON, M.A. Leaf appearance rate and final leaf number of switchgrass cultivars. **Crop Sci.**, v.37, p.864-870, 1997.

TAXA DE SENESCÊNCIA FOLIAR (TEMPO DE VIDA DA FOLHA)

- CARRERE et al. Tissue turnover within grass-clover mixed swards grazed by sheep. Methodology for calculating growth, senescence and intake fluxes. **J. Appl. Ecol.**, v.34, n.2, p.333-348, 1997.
- DURU, M., DUCROCQ, H. Growth and senescence of the successive leaves on a Cocksfoot tiller. Ontogenic development and effect of temperature. **Ann. Bot.**, v.85, p.635-643, 2000.
- DURU, M., DUCROCQ, H. Growth and senescence of the successive leaves on a Cocksfoot tiller. Effect of nitrogen and cutting regime. **Ann. Bot.**, v.85, p.645-653, 2000.
- GOMIDE, C.A.M., GOMIDE, J.A. Morfogênese de cultivares de *Panicum maximum* Jacq. **Rev. bras. zootec.**, v.29, n.2, p. 341-348, 2000.
- GRANT et al. Seasonal pattern of leaf growth and senescence of *Nardus stricta* and responses of tussocks to differing severity, timing and frequency of defoliation. **J. Appl. Ecol.**, v.33, n.5, p.1145-1155, 1996.
- HUNT, L.A. Some implications of death and decay in pasture production. **J. Br. Grassland Soc.**, v.20, n.1, p.27-31, 1965.
- MARRIOT, C.A. et al. Seasonal dynamics of leaf extension and losses to senescence and herbivory in extensively managed sown ryegrass-white clover swards. **J. Agric. Sci.**, v.132, n.1, p.77-89, 1999.

#### ALONGAMENTO DAS HASTES

- CÂNDIDO, M. J. D. ; GOMIDE, C. A. M. ; ALEXANDRINO, E. ; GOMIDE, J. A. ; PEREIRA, W. E. . Morfofisiologia do Dossel de *Panicum maximum* cv. Mombaça sob Lotação Intermitente com Três Períodos de Descanso. Revista Brasileira de Zootecnia-Brazilian Journal of Animal Science, Viçosa, v. 34, n. 2, p. 338-347, 2005.
- DAVIS, M.H., SIMMONS, S.R. Far-red light reflected from neighbouring vegetation promotes shoot elongation and accelerates flowering in spring barley plants. **Plant Cell Environ.**, v.17, n.7, p.829-836, 1994.

#### FOTOSSÍNTESE E RESPIRAÇÃO

- BRÉGARD, A., ALLARD, G. Sink to source transition in developing leaf blades of tall fescue. **New Phytol.**, v.141, n.1, p.45-50, 1999.
- HENNING, J.C. e BROWN, R.H. Effects of irradiance and temperature on photosynthesis in C<sub>3</sub>, C<sub>4</sub> and C<sub>3</sub>/C<sub>4</sub> *Panicum* species. **Photosynthesis Research**, 10: 101-112, 1986.
- HODGKINSON, K.C. The utilization of root organic compounds during the regeneration of lucerne. **Aust. J. Biol. Sci.**, v.22, p.1113-1123, 1969.
- HODGKINSON, K.C et al. The photosynthetic capacity of stubble leaves and their contribution to growth of the lucerne plant after high level cutting. **Aust. J. Agric. Res.** v.23, n.2, p.225-238, 1972.
- HODGKINSON, K.C. Influence of partial defoliation on photosynthesis, photorespiration and transpiration by lucerne leaves of different ages. **Aust. Plant Physiol.** v.1, n.4, p.561-578, 1974.
- KING, J.; SIM, E.M.; GRANT, S.A. Photosynthetic rate and carbon balance of grazed ryegrass pastures. **Grass Forage Sci.**, v.39, n.1, p.81-92, 1984.
- KING et al. Photosynthetic potential of ryegrass pastures when released from continuous stocking management. **Grass Forage Sci.**, v.43, n.1, p.41-48, 1988.
- LAMBERS, H. Does variation in photosynthetic rate explain variation in growth rate and yield?. **Neth. J. Agric. Sci.**, v.35, p.505-519, 1987.
- LAWLOR. Photosynthesis, productivity and environment. **J. Exper. Bot.**, v.46, p.1449-1461, 1995.

- PARSONS, A.J., LEAFE, E.F., COLLET, B., STILES, W. The physiology of grass production under grazing. I. Characteristics of leaf and canopy photosynthesis of continuously-grazed swards. **J. Appl. Ecol.**, v.20, n.1, p.117-126, 1983.
- PARSONS, A.J., LEAFE, E.L., COLLETT, B., PENNING, P.D., LEWIS, J. The physiology of grass production under grazing. II. Photosynthesis, crop growth and animal intake of continuously-grazed swards. **J. Appl. Ecol.**, v.20, n.1, p.127-139, 1983.
- PARSONS, A.J.; JOHNSON, I.R.; WILLIAMS, J.H.H. Leaf age structure and canopy photosynthesis in rotationally and continuously grazed swards. **Grass and Forage Science**, v.43, p.1-14, 1988.
- PERI, P.L., et al. Net photosynthetic rate of cocksfoot leaves under continuous and fluctuating shade conditions in the field. **Grass Forage Sci.**, v.57, p.157-170, 2002.
- ROBSON, M.J. The growth and development of simulated swards of perennial ryegrass. II. Carbon assimilation and respiration in a seedling sward. **Ann. Bot.**, v.37, n.151, p.501-518, 1973.
- RYLE, G.J.A., POWELL, C.E. Effect of rate of photosynthesis on the pattern of assimilate distribution in the graminaceous plant. **J. Exp. Bot.**, v.27, n.97, p.189-199, 1976.
- SINCLAIR, T.R., HORIE, T. Leaf nitrogen, photosynthesis, and crop radiation use efficiency: a review. **Crop Sci.**, v.29, n.1, p.90-98, 1989.
- WOLEDGE, J. The effect of light intensity during growth on the subsequent rate of photosynthesis of leaves of Tall Fescue (*Festuca arundinacea* Schreb). **Ann. Bot.**, v.35, n.140, p.311-322, 1971.
- WOLEDGE, J. The photosynthesis of ryegrass leaves grown in a simulated sward. **Ann. Appl. Biol.**, v.73, p.229-237, 1973.
- WOLEDGE, J. The effects of shading and cutting treatments on the photosynthetic rate of ryegrass leaves. **Ann. Bot.**, v.41, n.176, p.1279-1286, 1977.
- WOLEDGE, J. The effects of shading during vegetative and reproductive growth on the photosynthetic capacity of leaves in a grass sward. **Ann. Bot.**, v.42, n.181, p.1085-1089, 1978.
- WOLEDGE, J., LEAFE, E.L. Single leaf and canopy photosynthesis in a ryegrass sward. **Ann. Bot.**, v.40, p.773-783, 1976.
- ZELICH, I. The close relationship between net photosynthesis and crop yield. **Bioscience**, v.32, n.10, p.796-802, 1982.

#### ÍNDICE DE ÁREA FOLIAR

- BROWN, R.H., BLASER, R.E. Leaf area index in pasture growth. **Herbage Abst.**, v.38, n.1, p.1-8, 1968.
- DAVIDSON, J.L., MILTHORPE, F.L. Leaf growth in *Dactylis glomerata* following defoliation. **Ann. Bot.**, v.30, n.118, p.173-184, 1966.
- DAVIES, A. Leaf tissue remaining after cutting and regrowth in perennial ryegrass. **J. Agric. Sci.**, v.82, p.165-172, 1974.
- PEARCE et al. Relationships between leaf area index, light interception and net photosynthesis in orchardgrass. **Crop Sci.**, v.5, n.6, p.553-556, 1965.

#### RESERVAS ORGÂNICAS

- BOOYSEN e NELSON. Leaf area and carbohydrate reserves in regrowth of tall fescue. **Crop Sci.**, v.15, n.2, p.262-266, 1975.
- DAVIDSON, J.L., MILTHORPE, F.L. The effect of defoliation on the carbon balance in *Dactylis glomerata*. **Ann. Bot.**, v.30, n.118, p.185-198, 1966.

- De VISSER, R. et al. Kinetics and relative significance of remobilized and current C and N incorporation in leaf and root growth zones of *Lolium perenne* after defoliation: assessment by <sup>13</sup>C and <sup>15</sup>N steady-state labeling. **Plant Cell Environ.**, v.20, p.37-46, 1997.
- DONAGHY e FULKERSON. The importance of water-soluble carbohydrates reserves on regrowth and root growth of *Lolium perenne*. **Grass Forage Sci.**, v.52, p.401-407, 1997.
- DONAGHY e FULKERSON. Priority for allocation of water-soluble carbohydrate reserves during regrowth of *Lolium perenne*. **Grass Forage Sci.**, v.53, p.211-218, 1998.
- JOHANSSON, G. Carbon distribution in grass (*Festuca pratensis* L.) during regrowth after cutting – utilization of stored and newly assimilated carbon. **Plant & Soil**, v.151, n.1, p.11-20, 1993.
- MARSHALL, C., SGAR, G.R. The interdependence of tillers in *Lolium multiflorum* Lam. – a quantitative assessment. **J. Exp. Bot.**, v.19, n.61, p.785-794, 1968.
- MORVAN-BERTRAND, A., PAVIS, N., BOUCAUD, J., PRUD'HOMME, M.P. Partitioning of reserve and newly assimilated carbon in roots and leaf tissues of *Lolium perenne* during regrowth after defoliation: assessment by <sup>13</sup>C steady-state labeling and carbohydrate analysis. **Plant Cell Environ.**, v.22, p.1097-1108, 1999.
- PARSONS, A.J., ROBSON, M.J. Seasonal changes in the physiology of S24 perennial ryegrass (*Lolium perenne* L.). 4. Comparison of the carbon balance of the reproductive crop in spring and the vegetative crop in autumn. **Ann. Bot.**, v.50, n.\_, p.167-177, 1982.
- POWELL, J.M. et al. Nutrient and carbohydrate partitioning in Sorghum stover. **Agron. J.**, v.83, p.933-937, 1991.
- RICHARDS e CALDWELL. Soluble carbohydrates, concurrent photosynthesis and efficiency in regrowth following defoliation: a field study with *Agropyron* species. **J. Appl. Ecol.**, v.22, n.3, p.907-920, 1985.
- RYLE, G.J.A., POWELL, C.E. The export and distribution of <sup>14</sup>C-labelled assimilates from each leaf on the shoot of *Lolium temulentum* during reproductive and vegetative growth. **Ann. Bot.**, v.36, p.363-375, 1972.
- RYLE, G.J.A., POWELL, C.E. The utilization of recently assimilated carbon in graminaceous plants. **Ann. Appl. Biol.**, v.77, p.145-158, 1974.
- VOLENEC, J.J., NELSON, C.J. Carbohydrate metabolism in leaf meristems of Tall Fescue. **Plant Physiol.**, v.74, p.595-600, 1984.
- WARD e BLASER. Carbohydrate food reserves and leaf area in regrowth of orchardgrass. **Crop Sci.**, v.1, n.5, p.366-370, 1961.
- WATSON e WARD. Influence of intact tillers and height of cut on regrowth and carbohydrate reserves of dallisgrass (*Paspalum dilatatum* Poir.). **Crop Sci.**, v.10, n.5, p.474-476, 1970.
- WHITE, L.M. Carbohydrate reserves of grasses: a review. **J. Range Manage.**, v.26, n.1, p.13-18, 1973.

#### MERISTEMA APICAL

- BOTREL, M.A., GOMIDE, J.A. Importância do teor dos carboidratos de reserva e da sobrevivência dos meristemas apicais para a rebrota do capim-jaraguá (*Hyparrhenia rufa* (Ness) Stapf). **Rev. soc. bras. zootec.**, v.10, n.3, p. 411-426, 1981.

#### PERFILHAMENTO

- BEATY, E.R., et al. Yield, leaf growth, and tillering in bahiagrass by N rate and season. **Agron. J.**, v.69, p.308-311, 1977.
- BULLOCK et al. Tiller dynamics of two grasses - responses to grazing, density and weather. **J. Ecology**, v.82, n.2, p.331-340, 1994.
- CARVALHO, C.A.B. et al. Perfilhamento e acúmulo de forragem em pastagens de Florakirk (*Cynodon* spp.) sob pastejo. **B. indústr. anim.**, v.57, n.1, p.39-51, 2000.

- CARVALHO, C.A.B. et al. Demografia do perfilhamento e taxas de acúmulo de matéria seca em capim 'tifton 85' sob pastejo. **Scientia Agric.**, v.57, n.4, p. 591-600, 2000.
- CULVENOR, R.A. Observations on tillering in cultivars of phalaris under rotational grazing in a year with a summer-autumn drought. **Aust. J. Agric. Res.**, v.48, n.4, p.467-476, 1997.
- DAVIES, A., EVANS, M.E., EXLEY, J.K. Regrowth of perennial ryegrass as affected by simulated leaf sheaths. **J. Agric. Sci.**, v.101, p.131-137, 1983.
- DEREGIBUS, V.A. et al. Effects of light quality on tiller production in *Lolium* spp. **Plant Physiol.**, v.72, p.900-902, 1983.
- DEREGIBUS, V.A. et al. Tillering responses to enrichment of red light beneath the canopy in a humid natural grassland. **J. Appl. Ecol.**, v.22, p.199-206, 1985.
- GOMIDE, J.A., OBEID, J.A., RODRIGUES, L.R.A. Fatores morfofisiológicos de rebrota do capim-colônia *Panicum maximum*. **Rev. soc. bras. zootec.**, v.8, n.4, p. 532-562, 1979.
- HERNÁNDEZ GARAY, A. et al. Tiller size/density compensation in perennial ryegrass miniature swards subject to differing defoliation heights and a proposed productivity index. **Grass Forage Sci.**, v.54, n.4, p.347-356, 1999.
- HUME, D.E. Effect of cutting on production and tillering in prairie grass (*Bromus willdenowii* Kunth) compared with two ryegrass (*Lolium*) species. 1. Vegetative plants. **Ann. Bot.**, v.67, n.6, p.533-541, 1991.
- JEWISS, O.R. Tillering in grasses – its significance and control. **J. Br. Grassl. Soc.**, v.27, n.1, p.65-82, 1972.
- KLINK. Effects of clipping on size and tillering of native and African grasses of the Brazilian savannas (the cerrado). **Oikos**, v.70, p.365-376, 1994.
- LANGER, R.H.M. Tillering in herbage grasses. **Herbage Abst.**, v.33, n.3, p.141-148, 1963.
- LATERRA et al. Demography variability in tiller populations of two perennial pampa grasses. **J. Veget. Sci.**, v.8, p.369-376, 1997.
- LATERRA et al. Neighbour influence on the tiller demography of two perennial pampa grasses. **J. Veget. Sci.**, v.8, p.361-368, 1997.
- NASCIMENTO, M.P.S.C.B., NASCIMENTO, H.T., GOMIDE, J.A. Alguns aspectos morfofisiológicos de três gramíneas de clima tropical. **Rev. soc. bras. zootec.**, v.9, n.1, p. 142-158, 1980.
- NELSON, C.J., ZARROUGH, K.M. Tiller density and tiller weight as yield determinants of vegetative swards. In: OCCASIONAL SYMPOSIUM – PLANT PHYSIOLOGY AND HERBAGE PRODUCTION, 13, 1981, England. **Proceedings...** WRIGHT, C.E. (Ed.). England: British Grassland Society, 1981. p.25-29.
- PEDREIRA, J.V.S. Hábitos de perfilhamento do capim-colônia (*Panicum maximum*, Jacq.). **B. indústr. anim.**, v.32, n.1, p.111-114, 1975.
- RYLE, G.J.A. A comparison of leaf and tiller growth in seven perennial grasses as influenced by nitrogen and temperature. **J. Br. Grassl. Soc.**, v.19, n.3, p.281-290, 1964.
- SILSBURY, J.H. Interrelations in the growth and development of *Lolium* II. Tiller number and dry weight at low density. **Austr. J. Agric. Res.**, v.17, p.841-847, 1966.
- SIMON, J.C., LEMAIRE, G. Tillering and leaf area index in grasses in the vegetative phase. **Grass Forage Sci.**, v.42, p.373-380, 1987.
- SKINNER, R.H. e NELSON, C.J. Estimation of potential tiller production and site usage during tall fescue canopy development. **Ann. Bot.**, v.70, n.6, p.493-499, 1992.
- SKINNER, R.H. e NELSON, C.J. Effect of tiller trimming on phyllochron and tillering regulation during tall fescue development. **Crop Sci.**, v.34, n.5, p.1267-1273, 1994.
- VAN LOO, E.N. Tillering, leaf expansion and growth of plants of two cultivars or perennial ryegrass grown using hydroponics at two water potentials. **Ann. Botany**, v.70, n.6, p.511-518, 1992.
- WILLIAMS, R.D. Tillering in grasses cut for conservation, with special reference to perennial ryegrass. **Herbage Abst.**, v.40, n.4, p.383-388, 1970.

## FREQUENCIA E INTENSIDADE DE DESFOLHAÇÃO

- FULKERSON, W.J., SLACK, K. Leaf number as a criterion for determining defoliation time for *Lolium perenne*. 1. Effect of water-soluble carbohydrates and senescence. **Grass Forage Sci.**, v.49, n.4, p.373-377, 1994.
- FULKERSON, W.J., SLACK, K. Leaf number as a criterion for determining defoliation time for *Lolium perenne*. 2. Effect of defoliation frequency and height. **Grass Forage Sci.**, v.50, n.1, p.16-20, 1995.
- GOMIDE, C.A.M., GOMIDE, J.A., MARTINEZ Y HUAMAN, C.A. et al. Fotossíntese, reservas orgânicas e rebrota do Capim-Mombaça (*Panicum maximum* Jacq.) sob diferentes intensidades de desfolha do perfilho principal. **Rev. bras. zootec.**, v.31, n.6, p. 2165-2175, 2002.
- HODGSON, J. The frequency of defoliation of individual tillers in a set-stocked sward. **J. Br. Grassland Soc.**, v.21, n.4, p.258-263, 1966.
- HODGSON, J. e OLLERENSHAW, J.H. The frequency and severity of defoliation of individual tillers in set-stocked swards. **J. Br. Grassland Soc.**, v.24, n.3, p.226-234, 1969.
- KORTE, C.J., WATKIN, B.R., HARRIS, W. Use of residual leaf area index and light interception as criteria for spring-grazing management of ryegrass dominant pasture. **New Zealand J. Agric. Res.**, v.25, p.309-319, 1982.
- KORTE, C.J., WATKIN, B.R., HARRIS, W. Effects of the timing and intensity of spring-grazings on reproductive development, tillering, and herbage production of perennial ryegrass dominant pasture. **New Zealand J. Agric. Res.**, v.27, p.135-149, 1984.
- McKENZI, F. R. Influence of grazing frequency and intensity on the density and persistence of *Lolium perenne* tillers under subtropical conditions. **Tropical Grasslands.** v. 31, n. 3, p. 219-226, 1997.
- MELLO, A.C.L.; PEDREIRA, C.G.S. Respostas morfológicas do Capim-Tanzânia (*Panicum maximum* Jacq. cv. Tanzânia-1) irrigado à intensidade de desfolha sob lotação rotacionada. **Revista Brasileira de Zootecnia**, v.33, p. 282-289, 2004.
- PARSONS, A.J., PENNING, P.D. The effect of the duration of regrowth on photosynthesis, leaf death and the average rate of growth in a rotationally grazed sward. **Grass Forage Sci.**, v.43, n.1, p.15-27, 1988.
- VICKERY, P.J.; BRINK, V.C.; ORMROD, D.P. Net photosynthesis and leaf area index relationships in swards of *Dactylis glomerata* under contrasting defoliation regimes. **J. Br. Grassl. Soc.**, v.26, n.2, p.85-90, 1971.

## PRODUÇÃO E ACÚMULO DE FORRAGEM

- BIRCHAM, J.S. The effects of a change in herbage mass on herbage growth, senescence and net production rates in a continuously stocked mixed species sward. In: OCCASIONAL SYMPOSIUM – PLANT PHYSIOLOGY AND HERBAGE PRODUCTION, 13, 1981, England. **Proceedings...** WRIGHT, C.E. (Ed.). England: British Grassland Society, 1981. p.85-87.
- BIRCHAM, J.S. e HODGSON, J. The influence of sward condition on rates of herbage growth and senescence in mixed swards under continuous stocking management. **Grass Forage Sci.**, v.38, n.4, p.323-331, 1983.
- BIRCHAM, J.S., HODGSON, J. The effects of change in herbage mass on herbage growth and senescence in mixed swards. **Grass Forage Sci.**, v.39, n.2, p.111-115, 1984.
- CARVALHO, C.A.B., SILVA, S.C., SBRISSIA, A.F., FAGUNDES, J.L., CARNEVALLI, R.A., PINTO, L.F.M., PEDREIRA, C.G.S. Carboidratos não estruturais e acúmulo de forragem em pastagens de *Cynodon* spp. sob lotação contínua. **Scientia Agric.**, v.58, n.4, p. 667-674, 2001.

- FAGUNDES et al. Índice de área foliar, interceptação luminosa e acúmulo de forragem em pastagens de *Cynodon* spp. sob diferentes intensidades de pastejo. **Scientia Agric.**, v.56, n.4, p. 1141-1150, 1999.
- GRANT, S.A. et al. Sward management, lamina turnover and tiller population density in continuously stocked *Lolium perenne*-dominated swards. **Grass Forage Sci.**, v.38, n.4, p.333-344, 1983.
- HORST, G.L., NELSON, C.J., ASAY, K.H. Relationship of leaf elongation to forage yield of tall fescue genotypes. **Crop Sci.**, v.18, p.715-719, 1978.
- KORTE, C.J., WATKIN, B.R., HARRIS, W. Tillering in 'Grasslands nui' perennial ryegrass swards 1. Effect of cutting treatments on tiller appearance and longevity, relationship between tiller age and weight, and herbage production. **New Zealand J. Agric. Res.**, v.28, n.3, p.437-447, 1985.
- MAZZANTI, A., LEMAIRE, G., GASTAL, F. The effect of nitrogen fertilization upon the herbage production of tall fescue swards continuously grazed with sheep. 1. Herbage growth dynamics. **Grass Forage Sci.**, v.49, n.2, p.111-120, 1994.
- ROBSON, M.J. The growth and development of simulated swards of perennial ryegrass. I. Leaf growth and dry weight change as related to the ceiling yield of a seedling sward. **Ann. Bot.**, v.37, n.151, p.487-500, 1973.
- ZARROUGH et al. Relationship between tillering and forage yield of tall fescue. I. Yield. **Crop Sci.**, v.23, n.2, p.333-337, 1980.

#### ESTRUTURA DO DOSSEL

- BAKKER, M.L. et al. Effects of sward structure on the diet selected by guanacos (*Lama guanicoe*) and sheep (*Ovis aries*) grazing a perennial ryegrass-dominated sward. **Grass Forage Sci.**, v.53, n.1, p.19-30, 1998.
- BARTHAM, G.T. Sward structure and the depth of the grazed horizon. **Grass Forage Sci.**, v.36, n.2, p.130-131, 1981.
- BÉLANGER, G. Morphogenetic and structural characteristics of field-grown timothy cultivars differing in maturity. **Can. J. Plant Sci.**, v.76, n.2, p.277-282, 1996.
- BÉLANGER, G. Morphogenetic characteristics of timothy grown with varying N nutrition. **Can. J. Plant Sci.**, v.78, n.1, p.103-108, 1998.
- BURLISON, A.J.; HODGSON, J.; ILLIUS, A.W. Sward canopy structure and the bite dimensions and bite weight of grazing sheep. **Grass Forage Sci.**, v.46, n.1, p.29-38, 1991.
- CARRÈRE, P. et al. How does the vertical and horizontal structure of a perennial ryegrass and white clover sward influence grazing? **Grass Forage Sci.**, v.56, n.1, p.118-130, 2001.
- CHAPMAN, D.F., LEMAIRE, G. Morphogenetic and structural determinants of plant regrowth after defoliation. In: BAKER, M.J. ed. **Grasslands for our world**. Wellington: SIR Publishing, 1993. p.55-64.
- FAGUNDES, J.L., SIVA, S.C., PEDREIRA, C.G.S., CARNEVALLI, R.A., CARVALHO, C.A.B., SBRISSIA, A.F., PINTO, L.F.M. Intensidades de pastejo e a composição morfológica de pastos de *Cynodon* spp. **Scientia Agric.**, v.56, n.4, p. 897-908, 1999.
- FLORES, E.R.; LACA, E.; GRIGGS, T.C. et al. Sward height and vertical morphological differentiation determine cattle bite dimensions. **Agronomy Journal**, v.85, p.527-532, 1993.
- FORBES e COLEMAN. Influence of herbage mass and structure of warm-season grass on ingestive behavior of grazing cattle. In: INTERNATIONAL GRASSLAND CONGRESS, 16, 1985, Kyoto. **Proceedings...** Kyoto: Japanese Society of Grassland Science, 1985. p.1123-1125.
- GARCEZ NETO, A.F. et al. Respostas Morfogênicas e Estruturais de *Panicum maximum* cv. Mombaça sob diferentes níveis de adubação nitrogenada e alturas de corte. **Rev. bras. zootec.**, v.31, n.5, p. 1890-1900, 2002.

- GOMIDE, C.A.M.; GOMIDE, J.A.M. The duration of regrowth period and the structural traits in a rotationally grazed *Panicum maximum* sward. In: INTERNATIONAL GRASSLAND CONGRESS, 19, 2001, São Pedro. **Proceedings...** Piracicaba: FEALQ, 2001. (cd-rom).
- LACA, E.A., LEMAIRE, G. Measuring Sward Structure. In: MANNETJE, L., JONES, R.M. eds. **Field and Laboratory Methods for Grassland and Animal Production Research.** Wallingford: CAB International, 2000. p.103-121.
- LAIDLAW, A.S., WITHERS, J.A. Changes in contribution of white clover to canopy structure in perennial ryegrass/white clover swards in response to N fertilizer. **Grass Forage Sci.**, v.53, p.287-291, 1998.
- LANG, A.R.G. et al. Crop structure and the penetration of direct sunlight. **Agric. Forest Meteorol.**, v.35, p.83-101, 1985.
- McBEE e MILLER. Carbohydrate and lignin partitioning in sorghum stems and blades. **Agron. J.**, v.82, n. 3, p.687-690, 1990.
- MOORE et al. Canopy structure of *Aeschynomene Americana-Hemarthria altissima* pastures and ingestive behavior of cattle. In: INTERNATIONAL GRASSLAND CONGRESS, 16, 1985, Kyoto. **Proceedings...** Kyoto: Japanese Society of Grassland Science, 1985. p.1126-1128.
- OLIVEIRA et al. Características morfológicas e estruturais do capim-bermuda 'Tifton 85' (*Cynodon* spp) em diferentes idades de rebrota. **Rev. bras. zootec.**, v.29, n.6 (suplem.1), p. 1939-1948, 2000.
- RHODES, I. Productivity and canopy structure of two contrasting varieties of perennial ryegrass (*Lolium perenne* L.) grown in a controlled environment. **J. Br. Grassl. Soc.**, v.26, n.1, p.9-15, 1971.
- RHODES, I. The relationship between productivity and some components of canopy structure in ryegrass (*Lolium* spp.). **J. Agric. Sci.**, v.77, p.283-292, 1971.
- SHEEHY, J.E., PEACOCK, J.M. Microclimate, canopy structure and photosynthesis in canopies of three contrasting temperate forage grasses. **Ann. Bot.**, v.41, n.173, p.567-578, 1977.
- SILVA, D.S., GOMIDE, J.A., FONTES, C.A.A., QUEIROZ, A.C. Pressão de pastejo em pastagem de capim-elefante anão (*Pennisetum purpureum*, Schum. cv. Mott): 1 - Efeito sobre a estrutura e disponibilidade de pasto. **Rev. soc. bras. zootec.**, v.23, n.2, p.249-257, 1994.
- STOBBS, T.H. The effect of plant structure on the intake of tropical pastures. I. Variation in the bite size of grazing cattle. **Aust. J. Agric. Res.**, v.24, n.6, p.809-819, 1973.
- STOBBS, T.H. The effect of plant structure on the intake of tropical pastures. II. Differences in sward structure, nutritive value, and bite size of animals grazing *Setaria anceps* and *Chloris gayana* at various stages of growth. **Aust. J. Agric. Res.**, v.24, n.6, p.821-829, 1973.
- SUGIYAMA, S.; YONEYAMA, M.; TAKAHASHI, N. et al. Canopy structure and productivity of *Festuca arundinacea* Schreb. swards during vegetative and reproductive growth. **Grass and Forage Science**, v.40, p.49-55, 1985.
- UNGAR, E.D., NOY-MEIR, I. Herbage intake in relation to availability and sward structure: grazing processes and optimal foraging. **J. Appl. Ecol.**, v.25, n.3, p.1045-1062, 1988.
- UNGAR, E.D., RAVID, N. Bite horizons and dimensions for cattle grazing herbage to high levels of depletion. **Grass Forage Sci.**, v.54, n.4, p.357-364, 1999.
- UNGAR, E.D., RAVID, N., BRUCKENTAL, I. Bite dimensions for cattle grazing herbage at low levels of depletion. **Grass Forage Sci.**, v.56, p.35-45, 2001.

#### COMPORTAMENTO INGESTIVO

- DOUGHERTY, C.T. et al. Moderation of ingestive behaviour of beef cattle by grazing-induced changes in lucerne swards. **Grass Forage Sci.**, v.45, p.135-142, 1990.
- MOORE et al. Canopy structure of *Aeschynomene Americana-Hemarthria altissima* pastures and ingestive behavior of cattle. In: INTERNATIONAL GRASSLAND CONGRESS, 16, 1985, Kyoto. **Proceedings...** Kyoto: Japanese Society of Grassland Science, 1985. p.1126-1128.

## CONSUMO DE FORRAGEM

- ALLDEN, W.G., WHITTAKER, I.A.McD. The determinants of herbage intake by grazing sheep: the interrelationship of factors influencing herbage intake and availability. **Aust. J. Agric. Res.**, v.21, n.5, p.755-766, 1970.
- ALLEN, M.S. Effects of diet on short-term regulation of feed intake by lactating dairy cattle. **J. Dairy Sci.**, v.83, p.1598-1624, 2000.
- ARIAS et al. Structure of tall fescue swards and intake of grazing cattle. **Agron. J.**, v.82, n.3, p.545-548, 1990.
- BAKER, R.D., et al. The effect of herbage allowance upon the herbage intake and performance of suckler cows and calves. **Grass Forage Sci.**, v.36, n.3, p.189-199, 1981.
- BARRETT, P.D. et al. The effect of sward structure as influenced ryegrass genotype on bite dimensions and short-term intake rate by dairy cows. **Grass Forage Sci.**, v.58, n.1, p.2-11, 2003.
- BAUMONT, R. et al. How forage characteristics influence behaviour and intake in small ruminants: a review. **Livestock Production Science**, v.64, n.1, p.15-28, 2000.
- BOVAL, M., et al. The effect of herbage allowance on daily intake by Creole heifers tethered on natural *Dichanthium* spp. pasture. **Grass Forage Sci.**, v.55, p.201-208, 2000.
- BURNS, J. C.; POND, K.R.; FISHER, D.S. Measurement of forage intake. In: FAHEY JUNIOR, G.C. (Ed.). **Forage quality evaluation and utilization**. Madison: American Society of Agronomy, 1994. p.494-532.
- CAMPLING. Factors affecting the voluntary intake of grass. **J. Br. Grassland Soc.**, v.19, n.1, p.110-129, 1964.
- COMBELLAS, J., HODGSON, J. Herbage intake and milk production by grazing dairy cows. 1. The effects of variation in herbage mass and daily herbage allowance in a short-term trial. **Grass Forage Sci.**, v.34, p.209-214, 1979.
- COSGROVE, G.P. Grazing behaviour and forage intake. In: SIMPÓSIO INTERNACIONAL SOBRE PRODUÇÃO ANIMAL EM PASTEJO, 1997, Viçosa. **Anais...** GOMIDE, J.A. et al. (EE.). Viçosa: UFV, 1997. p. 59-80.
- DAVIES et al. Assessment of contrasting perennial ryegrasses, with and without white clover, under continuous sheep stocking in the uplands. 3. Herbage production, quality and intake. **Grass Forage Sci.**, v.46, n.1, p.39-49, 1991.
- DAVIES et al. Assessment of contrasting perennial ryegrasses, with and without white clover, under continuous sheep stocking in the uplands. 5. Herbage production, quality and intake in years 4-6. **Grass Forage Sci.**, v.48, n.3, p.213-222, 1993.
- DOUGHERTY, C.T., LAURIAULT, L. M., CORNELIUS, P.L., BRADLEY, N.W. Herbage allowance and intake of cattle. **J. Agric. Sci.**, v.112, p.395-401, 1989.
- DOUGHERTY, C.T., COLLINS, M., BRADLEY, N.W., CORNELIUS, P.L., LAURIAULT, L. M. Moderation of ingestive behaviour of beef cattle by grazing-induced changes in lucerne swards. **Grass Forage Sci.**, v.45, p.135-142, 1990.
- DOVE, H. Constraints to the modelling of diet selection and intake in the grazing ruminant. **Aust. J. Agric. Res.**, v.47, n.2, p.257-275, 1996.
- EDWARDS, G.R.; PARSONS, A.J.; PENNING, P.D. et al. Relationship between vegetation state and bite dimensions of sheep grazing contrasting plant species and its implications for intake rate and diet selection. **Grass Forage Sci.**, v.50, n.4, p.378-388, 1995.
- EUCLIDES, V.P.B., THIAGO, L.R.L.S., MACEDO, M.C.M., OLIVEIRA, M.P. Consumo voluntário de forragem de três cultivares de *Panicum maximum* sob pastejo. **Rev. bras. zootec.**, v.28, n.6, p. 1177-1185, 1999.
- EUCLIDES, V.P.B. et al. Consumo voluntário de *Brachiaria decumbens* cv. basilisk e *Brachiaria brizantha* cv. marandu sob pastejo. **Rev. bras. zootec.**, v.29, n.6, p. 2200-2208, 2000(suplem.2).

- FORBES, J.M. Integration of regulatory signals controlling forage intake in ruminants. **J. Anim. Sci.**, v.74, p.3029-3035, 1996.
- FORBES e COLEMAN. Influence of herbage mass and structure of warm-season grass on ingestive behavior of grazing cattle. In: INTERNATIONAL GRASSLAND CONGRESS, 16, 1985, Kyoto. **Proceedings...** Kyoto: Japanese Society of Grassland Science, 1985. p.1123-1125.
- FORBES, T.D.A., HODGSON, J. Comparative studies of the influence of sward conditions on the ingestive behaviour of cows and sheep. **Grass Forage Sci.**, v.40, n.1, p.69-77, 1985.
- GIBB e TREACHER. The effect of herbage allowance on herbage intake and performance of lambs grazing perennial ryegrass and red clover swards. **J. Agric. Sci.**, v.86, p.355-365, 1976.
- GIBB e TREACHER. The performance of weaned lambs offered diets containing different proportions of fresh perennial ryegrass and white clover. **Anim. Prod.**, v.39, n.3, p.413-420, 1984.
- GIBB, M.J., HUCKLE, C.A., NUTHALL, R., ROOK, A.J. Effect of sward surface height on intake and grazing behaviour by lactating Holstein Friesian cows. **Grass Forage Sci.**, v.52, p.309-321, 1997.
- HOWARD, M.D. et al. Voluntary intake and ingestive behavior of steers grazing Johnstone or endophyte-infected Kentucky-31 tall fescue. **J. Anim. Sci.**, v.70, n.4, p.1227-1237, 1992.
- KENNEY e BLACK. Factors affecting diet selection by sheep. I – Potential intake rate and acceptability of feed. **Aust. J. Agric. Res.**, v.35, n.4, p.551-563, 1984.
- LACA, E.A., WALLISDEVRIES, M.F. Acoustic measurement of intake and grazing behaviour of cattle. **Grass Forage Sci.**, v.55, p.97-104, 2000.
- LACA et al. An integrated methodology for studying short-term grazing behaviour of cattle. **Grass Forage Sci.**, v.47, n.1, p.81-90, 1992.
- LACA, E.A., UNGAR, E.D., SELIGMAN, N., DEMMENT, M.W. Effects of sward height and bulk density on bite dimensions of cattle grazing homogeneous swards. **Grass Forage Sci.**, v.47, n.1, p.91-102, 1992.
- MAYES, R.W. et al. The use of dosed and herbage n-alkanes as markers for the determination of herbage intake. **J. Agric. Sci.**, v.107, n.1, p.161-170, 1986.
- MAZZANTI, A., LEMAIRE, G. The effect of nitrogen fertilization upon the herbage production of tall fescue swards continuously grazed by sheep. 2. Consumption and efficiency of herbage utilization. **Grass Forage Sci.**, v.49, p.352-359, 1994.
- McCULLOUGH, M.E. The significance of and techniques used to measure forage intake and digestibility. **Agron. J.**, v.51, n.4, p.219-222, 1959.
- McGILLOWAY, D.A. et al. The relationship between level of sward height reduction in a rotationally grazed sward and short-term intake rates of dairy cows. **Grass Forage Sci.**, v.54, n.2, p.116-126, 1999.
- McMENIMAN, N.P. Methods of estimating intake of grazing animals. In: SIMPÓSIO INTERNACIONAL SOBRE TÓPICOS ESPECIAIS EM ZOOTECNIA/REUNIÃO DA SOCIEDADE BRASILEIRA DE ZOOTECNIA, 34, 1997, Juiz de Fora. **Anais...** Juiz de Fora: SBZ, 1997. (cd-rom)
- MERCHANT, M., RIACH, D.J. The intake and performance of cashmere goats grazing sown swards. **Grass Forage Sci.**, v.49, n.4, p.429-437, 1994.
- MOORE, J.E., SOLLENBERGER, E. Techniques to predict pasture intake. In: SIMPÓSIO INTERNACIONAL SOBRE PRODUÇÃO ANIMAL EM PASTEJO, 1997, Viçosa. **Anais...** GOMIDE, J.A. et al. (EE.). Viçosa: UFV, 1997. p. 59-80.
- PENNING, P.D., JOHNSON, R.H. The use of internal markers to estimate herbage digestibility and intake. 1. Potentially indigestible cellulose and acid insoluble ash. **J. Agric. Sci.**, v.100, n.1, p.127-131, 1983.
- PENNING, P.D., JOHNSON, R.H. The use of internal markers to estimate herbage digestibility and intake. 2. Indigestible acid detergent fibre. **J. Agric. Sci.**, v.100, n.1, p.133-138, 1983.

- PENNING, P.D., PARSONS, A.J., ORR, R.J., TREACHER, T.T. Intake and behaviour responses by sheep to changes in sward characteristics under continuous stocking. **Grass Forage Sci.**, v.46, n.1, p.15-28, 1991.
- PENNING, P.D., PARSONS, A.J., ORR, R.J., HOOPER, G.E. Intake and behaviour responses by sheep to changes in sward characteristics under rotational grazing. **Grass Forage Sci.**, v.49, n.4, p.476-486, 1994.
- PEYRAUD, J.L. et al. The effect of herbage allowance, herbage mass and animal factors upon herbage intake by grazing dairy cows. **Ann. Zootech.**, v.45, p.201-217, 1996.
- PRACHE, S. Intake rate, intake per bite and time per bite of lactating ewes on vegetative and reproductive swards. **Applied Animal Behaviour Science**, v.52, p.53-64, 1997.
- REEVES, M. et al. A comparison of three techniques to determine the herbage intake of dairy cows grazing kikuyu (*Pennisetum clandestinum*) pasture. **Aust. J. Exp. Agric.**, v.36, n.1, p.23-30, 1996.
- SMITH e REID. Use of chromic oxide as an indicator of fecal output for the purpose of determining the intake of pasture herbage by grazing cows. **J. Dairy Sci.**, v.38, n.5, p.515-524, 1955.
- SOLLENBERGER, L.E., BURNS, J.C. Canopy characteristics, ingestive behaviour and herbage intake in cultivated tropical grasslands. In: INTERNATIONAL GRASSLAND CONGRESS, 19, 2001, São Pedro. **Proceedings...** GOMIDE, J.A. (Ed.). Piracicaba: FEALQ, 2001. (compact disk).
- Van SOEST, P.J. Voluntary intake in relation to chemical composition and digestibility. **J. Anim. Sci.**, v.24, n.3, p.834-843, 1965.
- VAZQUEZ, O.P., SMITH, T.R. Factors affecting pasture intake and total dry matter intake in grazing dairy cows. **J. Dairy Sci.**, v.83, p.2301-2309, 2000.
- WADE, M.H., CARVALHO, P.C.F. Defoliation patterns and herbage intake on pastures. In: LEMAIRE, G. et al. (Eds.). **Grassland ecophysiology and grazing ecology**. Wallingford: CAB International, 2000. p.233-248.
- WESTON, R.H. Some aspects of constraint to forage consumption by ruminants. **Aust. J. Agric. Res.**, v.47, n.2, p.175-197, 1996.

#### QUALIDADE DA PASTAGEM

- DEL POZO, M. et al. Diet selection by sheep and goats and sward composition changes in a ryegrass/white clover sward previously grazed by cattle, sheep or goats. **Grass Forage Sci.**, v.52, p.278-290, 1997.
- ENGELS, F.M. e SCHUURMANS, J.L.L. Relationship between structural development of cell walls and degradation of tissues in maize stems. **J. Sci. Food Agric.** V.59(1), 45-51, 1992.
- SOLLENBERGER, L.E., JONES JUNIOR, C.S. Beef production from nitrogen-fertilized mott dwarf elephantgrass and Pensacola bahiagrass pastures. **Trop. Grassl.**, v.23, n.3, p.129-134, 1989.
- VAZQUEZ, O.P., SMITH, T.R. Factors affecting pasture intake and total dry matter intake in grazing dairy cows. **J. Dairy Sci.**, v.83, p.2301-2309, 2000.
- WILSON, J. R. Cell wall characteristics in relation to forage digestion by ruminants. **Journal of Agricultural Science**, v. 122, n. 2, p. 173-182, 1994.
- WILSON, J.R., KENNEDY, P.M. Plant and animal constraints to voluntary feed intake associated with fibre characteristics and particle breakdown and passage in ruminants. **Aust. J. Agric. Res.**, v.47, n.2, p.199-225, 1996.
- WILSON, J.R., MINSON, D.J. Prospects for improving the digestibility and intake of tropical grasses. **Trop. Grassl.**, v.14, n.3, p.253-259, 1980.
- WILSON, J. R.; MERTENS, D. R. Crop quality & utilization. **Crop Sci.**, v. 35, p. 251-259. 1995.
- WRIGHT, I.A. et al. Effects of grazing by sheep or cattle on sward structure and subsequent performance of weaned lambs. **Grass Forage Sci.**, v.56, p.138-150, 2001.

OFERTA DE FORRAGEM E EFICIÊNCIA DE USO DA FORRAGEM PRODUZIDA SOB  
PASTEJO

MAZZANTI, A., LEMAIRE, G. The effect of nitrogen fertilization upon the herbage production of tall fescue swards continuously grazed by sheep. 2. Consumption and efficiency of herbage utilization. **Grass Forage Sci.**, v.49, p.352-359, 1994.

STUTH, J.W.; KIRBY, D.R.; CHMIELEWSKY, R.E. Effect of herbage allowance on the efficiency of defoliation by the grazing animal. **Grass Forage Sci.**, v.36, n.1, p.9-15, 1981.

WILLIAMS, M.J., HAMMOND, A.C. Rotational vs. continuous intensive stocking management of Bahiagrass pasture for cows and calves. **Agron. J.**, v.91, n.1, p.11-16, 1999.